

## Exercise 06

## The Boltzmann distribution (2)

**Deadline:** Please hand in your protocol as a single pdf file to [saleksic@zedat.fu-berlin.de](mailto:saleksic@zedat.fu-berlin.de) by **Thursday, 23rd June, 10.15 a.m.** Protocol should contain Python code, plots, and comments.

**6.1 Microstates in a 5-level system (100 P)**

Consider a system with  $N$  particles and five different and equidistant energy levels (e.g.:  $\epsilon_1 = 0, \epsilon_2 = 1, \epsilon_3 = 2, \epsilon_4 = 3, \epsilon_5 = 4$ ). The configuration of the system at time  $t$  is defined by the energy levels of the individual particles. We can represent it as a vector, for example:

$$\mathbf{c}^\top(t) = (0, 4, 2, 1, 0, 3)$$

for a system with  $N = 6$  particles where  $\mathbf{c}^\top$  denotes the transpose of  $\mathbf{c}$ . We can change the configuration of the system, imposing the energy conservation, by making the following transitions

- Raise the energy of a randomly chosen particle  $i$  by 1
- lower the energy of a randomly chosen particle  $j$  by 1.

E.g., for  $i = 4$  and  $j = 2$  in the above example this yields

$$\mathbf{c}^\top(0) = (0, 4, 2, 1, 0, 3)$$

$$\mathbf{c}^\top(1) = (0, 3, 2, 2, 0, 3).$$

If  $i = j$ , the configuration does not change. However, we cannot increase the energy of a particle which is in the highest energy level, and we cannot decrease the energy of a particle which is in the lowest energy level.

- (a) Implement a program which generates a series of configurations according to this algorithm. In the initial state, all particles are in energy level  $\epsilon_2 = 1$ . (50 P)

To analyze your simulations,

- (c) Plot the total energy as a function of time. (10 P)
- (d) Plot the populations of the five energy levels as a function of time. (10 P)
- (e) Calculate the average population in each energy level and the standard deviation discarding the equilibration time from the trajectory. Plot them using the `errorbar(x,y,err)` function. (10 P)
- (f) Assuming that the particles are distributed according to the Boltzmann distribution, estimate  $\beta$ . (10 P)
- (g) Repeat the analysis with  $N = 10, 100, 1000, 10000$  particles for 10000 timesteps and discuss your results. (10 P)